



---

Presentation Abstract

[Add to Itinerary](#)

[Print](#)

---

Program#/Poster#: 71.11/P23

---

Presentation Title: Delay differential analysis: a framework for multimodal non-linear classification of Parkinson's disease

---

Location: Hall A

---

Presentation time: Saturday, Oct 17, 2015, 1:00 PM - 5:00 PM

---

Presenter at Poster: Sat, Oct. 17, 2015, 3:00 PM - 4:00 PM

---

Topic: ++D.16.c. Posture and gait: Higher order control, multi-task integration, and theory


---

Authors: **\*M. E. HERNANDEZ**<sup>1</sup>, J. WEYHENMEYER<sup>2</sup>, C. LAINSCSEK<sup>3</sup>, T. SEJNOWSKI<sup>3</sup>, H. POIZNER<sup>4</sup>;  
1Kinesiology and Community Hlth., UIUC, Urbana, IL;  
2Indiana Univ. Sch. of Med., Indianapolis, IN; 3The Salk Inst. for Biol. Studies, La Jolla, CA; 4Inst. for Neural Computation, UCSD, La Jolla, CA

---

Abstract: Parkinson's disease (PD) is the second most prevalent neurodegenerative disorder in the world, yet has no standard diagnostic test. PD is known to lead to marked alterations in cortico-thalamo-basal ganglia activity and subsequent movements, which may provide a biomarker for PD diagnosis. DDA is a time domain analysis

framework based on embedding theory in non-linear dynamics. An embedding reveals the nonlinear invariant properties of an unknown dynamical system (here the brain) from a single time series (EEG or behavioral signals). The DDA embedding serves as a low-dimensional nonlinear functional basis onto which the data are mapped. The combination of behavioral and neurological observations gives rise to a multimodal analysis framework that will improve the understanding and classification of neurological disease. We demonstrate how 750 ms of multimodal data can be used to improve DDA classification performance of PD after an unexpected perturbation of a virtual target during reach to grasp movements. We found that the anteroposterior hand position and hand aperture, in particular, provide improved classification performance in comparison to clean EEG data, as evaluated by the area under the ROC curve (AROC), (AROC increases from 0.71 to 0.81 with the addition of behavioral data). Thus, multimodal DDA may provide a tool for aiding the clinician in the diagnosis of PD and allow for earlier intervention with disease modifying therapeutics.



Disclosures: **M.E. Hernandez:** None. **J. Weyhenmeyer:** None. **C. Lainscsek:** None. **T. Sejnowski:** None. **H. Poizner:** None.

Keyword (s): PARKINSON'S DISEASE

NEUROIMAGING

MOTOR CONTROL

Support: Howard Hughes Medical Institute

---

NSF grant #SMA-1041755

---

NSF ENG-1137279 (EFRI M3C)

---

ONR MURI Award No.: N00014-10-1-0072

---

**Note:** When adding items to your Itinerary, please click "Add Checked Selections to My Itinerary" on EACH page of your search results.

[At the Meeting](#)

[Sessions/Events](#)

[Abstracts](#)

[Registration](#)

[Hotel/Travel](#)

[Exhibits](#)

[Fellowships, Awards, and Prizes](#)

[Frequently Asked Questions](#)

[CME](#)

[Access the SFN Member Center](#)

OASIS Technical Support.  
Monday - Friday, 9 am - 5 pm CT  
Phone: 1-217-398-1792  
Email: [OASIS Helpdesk](#)

---

[Leave OASIS Feedback](#)

---

The Online

Powered by



Abstract  
Submission and  
Invitation  
System  
© 1996 - 2015  
Coe-Truman  
Technologies,  
Inc. All rights  
reserved.

Services by



Coe-Truman Technologies, Inc.