

## Mini-review

## Does schizophrenia arise from oxidative dysregulation of parvalbumin-interneurons in the developing cortex?

M. Margarita Behrens<sup>a,\*</sup>, Terrence J. Sejnowski<sup>a,b</sup><sup>a</sup>Howard Hughes Medical Institute, The Salk Institute for Biological Studies, La Jolla, CA 92037, USA<sup>b</sup>Division of Biological Sciences, University of California at San Diego, La Jolla, CA, 92093, USA

## ARTICLE INFO

## Article history:

Received 14 April 2009

Received in revised form

30 May 2009

Accepted 1 June 2009

## Keywords:

Redox

Parvalbumin

Fast-spiking

Gamma oscillations

GABAergic

Interleukin-6

NADPH oxidase

Schizophrenia

## ABSTRACT

An imbalance in the redox-state of the brain may be part of the underlying pathophysiology in schizophrenia. Inflammatory mediators, such as IL-6, which can tip the redox balance into a pro-oxidant state, have been consistently found to be altered in schizophrenia patients. However, the relationship of altered redox-state to altered brain functions observed in the disease has been unclear. Recent data from a pharmacological model of schizophrenia suggest that redox and inflammatory imbalances may be directly linked to the pathophysiology of the disease by alterations in fast-spiking interneurons. Repetitive adult exposure to the NMDA-R antagonist ketamine increases the levels of the proinflammatory cytokine interleukin-6 in brain which, through activation of the superoxide-producing enzyme NADPH oxidase (Nox2), leads to the loss of the GABAergic phenotype of PV-interneurons and to decreased inhibitory activity in prefrontal cortex. This effect is not observed after a single exposure to ketamine, suggesting that the first exposure to the NMDA-R antagonist primes the brain such that deleterious effects on PV-interneurons appear upon repetitive exposures. The effects of activation of the IL-6/Nox2 pathway on the PV-interneuronal system are reversible in the adult brain, but permanent in the developing cortex. The slow development of PV-interneurons, although essential for shaping of neuronal circuits during postnatal brain development, increases their vulnerability to deleterious insults that can permanently affect their maturational process. Thus, in individuals with genetic predisposition, the persistent activation of the IL-6/Nox2 pathway may be an environmental factor that tips the redox balance leading to schizophrenia symptoms in late adolescence and early adulthood.

© 2009 Elsevier Ltd. All rights reserved.

## 1. Introduction

The pathophysiology of schizophrenia is complex and involves many different cortical and subcortical systems. In particular, fast-spiking parvalbumin (PV)-positive inhibitory neurons, which represent 5% of all cortical neurons, are strongly affected. Reduced expression of GAD67, the main isoform synthesizing GABA in brain, is one of the most replicated findings in schizophrenia postmortem brain studies (Benes and Berretta, 2001; Lewis et al., 2005), and single nucleotide polymorphisms in the regulatory region of *Gad1* (the gene coding for GAD67) are associated with childhood onset of schizophrenia (Rapoport et al., 2005). The decrease in GAD67 occurs primarily in the subset of inhibitory interneurons expressing the calcium binding protein parvalbumin (Beasley and Reynolds, 1997; Hashimoto et al., 2003). This apparent “loss of GABAergic

phenotype” in PV-interneurons led to the suggestion that dysfunction of these fast-spiking inhibitory interneurons may be a core feature of the disease (Lewis et al., 2005). Whether these deficiencies are a consequence or a cause of the disorder is, however, a matter of debate.

PV-interneurons are involved in the generation of gamma oscillations, which regulate working memory and information transmission between cortical areas (Salinas and Sejnowski, 2001; Bartos et al., 2007). In particular, synaptic inhibition from PV-interneurons controls the firing rates of pyramidal neurons, synchronizes spikes within populations of neurons, and participates in the development of executive functions associated with prefrontal brain regions (Kawaguchi and Kubota, 1993; Goldman-Rakic, 1999; Markram et al., 2004). PV-interneurons are a part of the network that generates oscillatory activity in the gamma range (Sohal et al., 2009; Cardin et al., 2009), suggesting that their dysfunction may account for the disruption in evoked gamma-frequency oscillations and as well as the cognitive deficits observed in schizophrenia (Gonzalez-Burgos and Lewis, 2008; Roopun et al., 2008; Uhlhaas et al., 2008).

\* Corresponding author. Tel.: +1 858 453 4100.

E-mail address: [mbehrens@salk.edu](mailto:mbehrens@salk.edu) (M.M. Behrens).

Adult levels of executive function emerge relatively late in the postnatal development of primates (Alexander and Goldman, 1978) and probably of rodents (Bachevalier and Beauregard, 1993; Ba and Seri, 1995). In primates and rodents, the delay in achieving mature performance on executive function tasks correlates with the maturation of oscillatory activity in the gamma range and with the maturation of PV-interneuronal networks (Wilson et al., 1994; Rao et al., 2000; Doischer et al., 2008; Uhlhaas et al., 2009), consistent with the delayed maturation of PV-inhibitory circuits. Development of PV-synaptic contacts, which sculpt this inhibitory network throughout childhood and adolescence, is dependent on GAD67 synthesis and activity in rodents (Chattopadhyaya et al., 2007). Environmental insults affecting the development of this inhibitory network, e.g. by affecting GAD67 expression, may lead to the abnormal formation of synaptic contacts by these interneurons. Thus, the dependence on GAD67 expression could make the developing brain vulnerable to environmental inputs that through disruption of the normal development of this inhibitory circuit lead to psychiatric diseases in adulthood.

## 2. The NMDA receptor antagonist model of schizophrenia

Several animal models recapitulating aspects (endophenotypes) of schizophrenia have been developed. Among these, exposure to NMDA receptor (NMDA-R) antagonists such as phencyclidine (PCP), ketamine, and MK801 are widely used in adult animals as acute pharmacological models to study behavioral and neurochemical disruptions relevant to the disease (rev. in Mouri et al., 2007). Administration of PCP to rodents produces deficits in spatial working memory, in reversal learning, and in sustained attention (Jentsch and Roth, 1999; Stefani and Moghaddam, 2005b). Similarly, ketamine can impair performance on tasks testing executive function in humans (Krystal et al., 2005) and non-human primates (Stoet and Snyder, 2006), resembling executive functioning deficits that are associated with treatment-refractory aspects of schizophrenia (Kerns et al., 2008).

At the neurochemical level, acute exposure to NMDA-R antagonists in adulthood increases excitatory transmission in frontal and anterior cingulate cortex across species (Takahata and Moghaddam, 2003). This hyper-excitation has been related to an increase in thalamo-cortical glutamatergic excitation of downstream cortical regions such as the anterior cingulate and retrosplenial cortices (Tomitaka et al., 2000; Holcomb et al., 2005). PET scan studies have shown that schizophrenic subjects respond to ketamine with higher hypermetabolism than normal subjects. Even under resting conditions schizophrenia patients show altered activity in the prefrontal cortex (PFC) and parahippocampal regions (Garrity et al., 2007) and suffer from persistent symptoms and chronic deficiency in their cognitive ability.

Although acute exposures to NMDA-R antagonists can produce some symptoms of schizophrenia in healthy subjects, these are transient and disappear after washout of the drug. Repetitive NMDA-R antagonist treatment in animals produce more persistent effects on stereotypy and locomotor activity, as well as enduring cognitive deficits and neurochemical changes that resemble more accurately the alterations observed in schizophrenia (Jentsch and Roth, 1999; Morris et al., 2005; Mouri et al., 2007). For example, the initial hypermetabolism, observed after acute NMDA-R antagonist exposure, is followed by a decrease in metabolic activity in the PFC, as well as within structures of the auditory system, and the reticular nucleus of the thalamus (Cochran et al., 2003). Repetitive PCP decreases dopamine in the dorsolateral prefrontal cortex, prelimbic cortex, and cingulate cortex, but not in supplementary motor area (Jentsch and Roth, 1999). This regimen also elicits alterations in N-acetylaspartate (NAA) and N-acetylaspartylglutamate (NAAG) in

temporal cortex and hippocampus (Reynolds et al., 2005), and decreases 5HT<sub>2A</sub> receptor binding in the PFC (Steward et al., 2004) in close similarity to schizophrenia pathology (Laruelle et al., 1993; Nudmamud et al., 2003). Furthermore, while the behavioral and neurochemical effects of acute exposures to NMDA-R antagonists are believed to occur upon disinhibition of cortical excitatory activity due to increased sensitivity of inhibitory systems to blockade of NMDA-R function (Olney et al., 1999; Homayoun and Moghaddam, 2007; Lisman et al., 2008; Middleton et al., 2008), they do not lead to the enduring changes in PV-interneurons observed in schizophrenia. Repetitive exposures to NMDA-R antagonists, on the other hand, produce a reduction in GAD67 expression in PV-interneurons of rodents (Behrens et al., 2007, 2008) as well as decreased expression of parvalbumin in rodents and non-human primates (Cochran et al., 2002; Keilhoff et al., 2004; Rujescu et al., 2006; Morrow et al., 2007).

## 3. The role of superoxide in the persistent effects of NMDA-R antagonists

Although exposure to one injection of the NMDA-R antagonist ketamine does not lead to the loss of GABAergic phenotype of PV-interneurons, injections of ketamine on two consecutive days is sufficient to produce this loss (Behrens et al., 2008). Similar exposures to ketamine in rats lead to an enduring decrease of inhibitory tone in prefrontal cortex (Zhang et al., 2008), supporting the idea that repetitive exposures to NMDA-R antagonists produce enduring effects resembling those observed in schizophrenia.

The repetitive, but not acute, exposure to NMDA-R antagonists disinhibited excitatory circuits and activated the superoxide-producing enzyme NADPH oxidase-2 (Nox2) in three week old primary cortical neurons as well as in adult brain. Furthermore, the inhibition of Nox2 with apocynin or eliminating superoxide with a brain-penetrant SOD-mimetic prevented the loss of phenotype of PV-interneurons *in vitro* and *in vivo*, and the ketamine effects were absent in Nox2-deficient animals (Behrens et al., 2007, 2008). Exposure to PCP and more selective NMDA-R antagonists such as MK801 and CPP were shown to produce a rapid increase in reactive oxygen- and nitrogen-species (ROS) *in vitro* (Xia et al., 2002), and *in vivo* (Zuo et al., 2007; Fejgin et al., 2008) and repetitive exposures *in vivo* led to a substantial elevation of baseline levels of free radicals, suggesting that this treatment results in a persistent change in the oxidative state of the cortex (Zuo et al., 2007). Interestingly, recent results have shown that NMDA receptor activity is required for the expression of antioxidant enzymes (Papadia et al., 2008), further supporting the idea that prolonged blockade of NMDA receptors produces an increased oxidative state.

Several studies have reported increased oxidative and nitrosative state, as well as a diminished antioxidant capacity in schizophrenia patients (reviewed in Do et al., 2009). Glutathione (GSH), responsible for detoxification of reactive oxygen and other radical species, is consistently decreased in cerebrospinal fluid of drug-naïve schizophrenia patients (Do et al., 2000), as well as in postmortem tissue (Yao et al., 2006). Polymorphisms in genes coding for enzymes that participate in GSH synthesis have been linked to schizophrenia risk (Tosic et al., 2006; Gysin et al., 2007), and acute frontal-brain GSH depletion in adult rodents was recently shown to produce disruptions in short-term memory, supporting the link between depletion of brain GSH levels and cognitive impairments that occur in schizophrenia (Pileblad et al., 1989; Jacobsen et al., 2005; Dean et al., 2009). Moreover, acute GSH depletion potentiates the release of dopamine produced by amphetamine in striatum and potentiates the behavioral effects of NMDA-R antagonists as well as those of amphetamine (Jacobsen et al., 2005). Cysteine, the limiting substrate in the synthesis GSH in

neurons, is transported from the extracellular space by the main glutamate transporter present in neurons, EAAC1 (Aoyama et al., 2006). EAAC1 as well as EAAC2 have been shown to be highly sensitive to oxidative conditions: reducing agents activate, and oxidation inactivates glutamate transport (Trotti et al., 1997). Thus, the increased superoxide production caused by activation of the IL-6/Nox2 pathway after exposure to NMDA-R antagonists would be expected to produce the redox inactivation of EAAC1 and a decrease in cysteine transport, leading to diminished GSH content in brain. If such activation of the IL-6/Nox2 pathway occurs in situations of genetically diminished brain GSH levels, such as those described in some schizophrenia cohorts, it will lead to further decrease in GSH levels, NMDA-R hypofunction and increased oxidative damage to macromolecules and lipids (Do et al., 2009). Recent results showing that treatment with N-acetyl-cysteine, a precursor of GSH, improves negative symptoms, and corrects mismatch negativity in schizophrenia patients further support the idea of a redox imbalance in schizophrenia (Berk et al., 2008; Lavoie et al., 2008).

#### 4. Mechanism of activation of Nox2 in neurons

One of the most consistent findings in schizophrenia patients is an imbalance in plasma and cerebrospinal fluid levels of cytokines (Muller et al., 2000). In particular, elevated plasma levels of IL-6 have been consistently reported in patients and first-degree relatives with mood disorders (Ganguli et al., 1994; Naudin et al., 1996; Nunes et al., 2005), and correlate with exacerbation of psychotic episodes (Ganguli et al., 1994; Naudin et al., 1996; Lin et al., 1998; Zhang et al., 2002; Kudoh et al., 2003; Nunes et al., 2005). Furthermore, treatment with atypical antipsychotics can alter circulating cytokines (Pollmacher et al., 2000). The increased levels of cytokines in schizophrenia patients, together with the important role played by Nox2-dependent NADPH oxidase in inflammatory processes outside the CNS suggests the possible involvement of proinflammatory molecules in the effects of NMDA-R antagonists.

Prolonged exposure to ketamine *in vitro* or repetitive exposures *in vivo* increased the levels of IL-6 mRNA, without affecting the levels of IL-1 $\beta$  or TNF $\alpha$  mRNAs (Behrens et al., 2008). Furthermore, brain IL-6 production is necessary and sufficient to produce the induction and activation of Nox2, and the consequent loss of the GABAergic phenotype of PV-interneurons (Behrens et al., 2008). Interestingly, increased activity of Nox2 was shown in neutrophils isolated from schizophrenia patients, which correlated with negative symptoms (Sirota et al., 2003), suggesting that the increased IL-6 levels in schizophrenia patients may also lead to induction of the peripheral enzyme.

However, although the effects of repetitive exposures to NMDA-R antagonist resemble more accurately the dysfunction of PV-interneurons observed in schizophrenia patients, these effects are slowly reversible in both the *in vitro* and *in vivo* models, requiring 48 h in culture (Kinney et al., 2006) or several days in the absence of drug *in vivo* (Behrens et al., 2008). Thus, although causing persistent effects, repetitive adult exposures to NMDA-R antagonists do not produce the irreversible changes in inhibitory circuitry observed in schizophrenia.

#### 5. Neurodevelopmental origins of schizophrenia: activation of the IL-6/Nox2 pathway may alter the development of PV-interneurons

Mild developmental impairments caused either by a genetic predisposition or by immune activation during development are believed to contribute to the appearance of schizophrenic symptoms in early adulthood (Rapoport et al., 2005). There is a strong correlation between infections in mid-gestation and the incidence

of schizophrenia in the offspring (reviewed in Brown, 2006; Paterson, 2008). Cytokine induction due to abnormal immune activation, or inflammation, derails normal brain development leading to alterations in cognition in adulthood (Gilmore and Jarskog, 1997; Nawa et al., 2000; Smith et al., 2007). Recently, using a rodent maternal-infection model Smith and collaborators found that maternal IL-6 induction during infection is responsible for the delayed schizophrenia-like behavior observed in the adult offspring (Smith et al., 2007). A lasting imbalance in cytokine levels was observed in the offspring throughout the first month of age. Since activation of the IL-6/Nox2 pathway and consequent increase in superoxide production in brain is required for the reversible loss of phenotype of PV-interneurons in adulthood (Behrens et al., 2008), activation of this pathway may be responsible for the delayed effects observed in the maternal-infection model of schizophrenia. Furthermore, decreased antioxidant capacity during early postnatal period induce cognitive derangements relevant to schizophrenia, as well as a decreased number of PV-interneurons in adulthood (Cabungcal et al., 2007).

Several neurodevelopmental models of schizophrenia converge on a sustained dysfunction of the PV-interneuronal system occurring during late prenatal/early postnatal development. Studies in the maternal-infection model, the DISC1 model, and prenatal exposure to methylazoxymethanol acetate (MAM), have shown that alterations of brain development during specific periods of pre or postnatal development produce discrete disruptions that lead to behavioral and neurochemical effects that include a decreased number of PV-interneurons (Hikida et al., 2007; Lodge and Grace, 2007; Meyer et al., 2007; Lodge et al., 2009). In the MAM model, where the mitotoxin is applied during interneuronal proliferation/migration stage, the number of PV-interneurons decreased in specific brain regions that correlated with alterations in oscillatory activity and decreased lateral inhibition in adulthood (Lodge et al., 2009).

In rodents and primates, maturation of the PV-interneuronal system occurs postnatally (reviewed in Lewis et al., 2004; Huang, 2009). Studies performed in rodents show that migration from the medial ganglionic eminence to the cortical plate is complete by around embryonic day 15, but the neurons remain silent until the beginning of the second postnatal week when their maturation begins. Before the first week of age, parvalbumin expression is absent in these neurons, but appears when volleys of excitatory activity arrive from thalamo-cortical projections during the second postnatal week. Further maturation of PV-interneuron synaptic contacts develops independently of thalamic inputs and reaches adult levels by the end of the first month of age (Di Cristo et al., 2004). A recent study in mouse cortex showed pronounced transcriptional and electrophysiological changes in PV-interneurons occurring during the first month of postnatal development (Okaty et al., 2009). Profound changes in the ion-channel repertoire expressed at different time-points during the first postnatal month shape the electrophysiological maturation required to attain the characteristic fast-spiking non-accommodating current patterns observed in adulthood. Oscillatory activity in the gamma range also begins during this second postnatal week, and matures throughout the following weeks to reach adult levels by the end of the adolescent period (Doischer et al., 2008). In non-human primates, the appearance of parvalbumin expression and development of PV-synaptic contacts begins at around 3 months of age, and develops profusely throughout childhood and adolescence (Reynolds and Beasley, 2001; Cruz et al., 2003). During adolescence, however, the restructuring and pruning of synaptic contacts reduces the levels to those found in adulthood (Cruz et al., 2003). Cortical neural synchrony and cognitive performance also show a protracted maturation in humans, reaching mature levels only in early adulthood after a period of decreased cognitive performance

and synchrony during adolescence (Uhlhaas et al., 2009). The profound reorganization of synaptic activity that occurs during adolescence could be responsible for the significant reductions in phase synchronization observed. The emergence of psychotic symptoms in schizophrenia during late adolescence/early adulthood could be a consequence of this reorganization process in compromised inhibitory networks (Feinberg, 1982).

The pro-psychotic effects of NMDA-R antagonists appear only in early adulthood and exposure of four week old rats to ketamine on two consecutive days did not reduce GAD67 immunoreactivity (Zhang et al., 2008). However, 24 h continuous exposure of three-week-old cultured neurons to ketamine produces a substantial reduction in GAD67 immunoreactivity (Kinney et al., 2006). These results suggest that PV-interneurons are more resistant to the effects of NMDA-R antagonists during the rodent equivalent to the adolescence period in humans. The overwhelming proliferation of PV-synaptic contacts observed before adolescence (Cruz et al., 2003), as well as the slow functional maturation of PV-inhibitory network (Doischer et al., 2008; Okaty et al., 2009) may explain the lack of pro-psychotic effects of NMDA-R antagonists before early adulthood. However, hormonal influences or changes in receptor-expression patterns (Okaty et al., 2009) and changes in response to modulatory neurotransmitters (Tseng and O'Donnell, 2007) that occur during adolescence could also explain the lack of pro-psychotic effects of NMDA-R antagonists.

In contrast to the reversible effects observed in adult animals (Behrens et al., 2008), embryonic and repetitive exposures during the second postnatal week to NMDA-R antagonists can produce loss of PV-interneurons and persistent behavioral and neurochemical deficits that appear only in adulthood (Stefani and Moghaddam, 2005a; Mouri et al., 2007; Wang et al., 2008). Mice expressing 10% of the normal NMDA-R activity develop cognitive disruptions resembling those found in schizophrenia (reviewed in (Gainetdinov et al., 2001)). Ablation of NR1 subunits in GABAergic neurons during early postnatal development leads to a loss of parvalbumin expression in PV-interneurons in adulthood and to behavioral disruptions reminiscent to those found in schizophrenia (Belforte et al., 2008). Taken together, these results support the idea of a critical role for NMDA-R function in the postnatal maturation of PV-interneurons, and raise a note of caution in the use of anesthetics with demonstrated NMDA-R antagonist activity in children.

Ketamine, although not approved for use in humans less than 17 years of age, is commonly used as an anesthetic in children (Mellon et al., 2007). Ketamine and other NMDA-R antagonists cause neurodegeneration in the developing brain in rodents and primates (Olney, 2002; Wang and Slikker, 2008). Although the doses used in these experiments are higher relative to those used in humans, repetitive sub-anesthetic doses, by activating the IL-6/Nox2 pathway in brain, may halt the maturation process of PV-interneurons and lead to a permanent dysfunction of cognitive processes. This is supported by the finding that antioxidants applied during perinatal exposure to NMDA-R antagonists prevent the development of behavioral disruptions (Wang et al., 2003), and that Nox2-deficient mice are protected from the decrease in PV-interneurons observed in animals that were exposed to NMDA-R antagonists during the second postnatal week (Shehktman and Behrens, unpublished).

## 6. Glutamate receptors and the GABAergic phenotype of PV-Interneurons

In common with other glutamatergic synapses, glutamate preferentially activates AMPA-type glutamate receptors in mature PV-interneurons. However, unlike receptors on pyramidal neurons, AMPA receptors on PV-interneurons do not express GluR2 subunits,

making them highly  $\text{Ca}^{2+}$  permeable (Goldberg et al., 2003). Regarding group I metabotropic glutamate receptors, cortical and hippocampal PV-interneurons preferentially express mGluR5 (Cauli et al., 1997; van Hooft et al., 2000). On the other hand, NMDA-Rs in PV-interneurons have a subunit composition that differs from neighboring pyramidal neurons, with NR2A and NR2C subunits being highly expressed (Kinney et al., 2006; Xi et al., 2009). NMDA-Rs exert a tight control of the basal excitability in PV-interneurons, and are highly sensitive to NMDA-R antagonists (Grunze et al., 1996; Li et al., 2002; Middleton et al., 2008). Given this composition of glutamate receptors, it is expected that glutamatergic transmission in PV-interneurons will be substantially different than transmission in pyramidal neurons.

Blockade of NMDA-Rs during the third week of development in culture leads to the loss of the GABAergic phenotype of PV-interneurons, and this loss can be prevented by strategies that increase intracellular calcium levels (Kinney et al., 2006). As noted above, PV-interneurons express GluR2-less AMPA receptors and are permeable to  $\text{Ca}^{2+}$ . Since the initial disinhibition caused by the NMDA-R antagonists lead to an increase in excitatory transmission, it would be expected that AMPA receptors in PV-interneurons would permeate enough  $\text{Ca}^{2+}$  to prevent the deleterious effects of blockade of NMDA-Rs. Furthermore, the increase in excitatory transmission caused by disinhibition should activate group I metabotropic glutamate receptors present in PV-interneurons and also lead to increases in intracellular  $\text{Ca}^{2+}$ , now through release from intracellular stores. However, co-exposure to a calcium-channel opener or an activator of group I metabotropic receptors was needed to preserve the GABAergic phenotype of PV-interneurons in the presence of NMDA-R antagonists (Kinney et al., 2006). This lack of response of AMPA and mGluR5 receptors to the increased excitatory transmission after disinhibition led to the hypothesis that prolonged blockade of NMDA-Rs in PV-interneurons results in an enduring change in AMPA and mGluR5-mediated responses to glutamate (Behrens et al., 2007).

What is the difference between PV-interneurons and pyramidal neurons that makes the former so sensitive to NMDA-R antagonists? Perhaps the specific subunit composition of the glutamate receptor in PV-interneurons is responsible. NMDA-Rs containing NR2C subunits have a reduced  $\text{Mg}^{2+}$  block, and consequently may be highly sensitive to ambient glutamate concentrations. Under normal physiological conditions  $\text{Mg}^{2+}$  concentrations are high enough to block NR2A/B-containing receptors, whereas NR2C and NR2D containing receptors will not show such blockade. Recent evidence shows that at physiological  $\text{Mg}^{2+}$  concentrations NMDA-R antagonists such as ketamine or memantine have negligible inhibitory effects on NR2A or NR2B containing receptors, whereas their inhibitory effects on NR2C or NR2D containing receptors remain unaltered (Kotermanski and Johnson, 2009). Thus, since NMDA-Rs in PV-interneurons contain higher levels of NR2A and NR2C receptors, antagonists such as ketamine would have stronger effects in these interneurons than on pyramidal neurons. In the presence of NMDA-R antagonists, lack of function of NR2A/NR2C containing receptors in PV-interneurons may then lead to profound changes in the function of this inhibitory system. Indeed, repetitive exposure to MK801 in vivo was recently shown to decrease the expression of NMDA-R subunits in PV-interneurons in the prefrontal cortex of rats (Xi et al., 2009). Interestingly, the NR2C subunit decreased most (~87-fold) after treatment.


## 7. Redox dysregulation of NMDA-R mediated transmission in PV-interneurons

Inactivation of synaptic proteins through oxidation is a well-described phenomenon, and considered to be behind many of the

derangements of the nervous system observed in disease states (Rowan et al., 2005; Butterfield, 2006; Satoh and Lipton, 2007). Regulatory redox sites have been found in many proteins that are key to glutamatergic neurotransmission including, serine-race-mase that is responsible for the synthesis of the endogenous modulator of the glycine site in NMDA receptors (Mustafa et al., 2007); glutamine synthase that is responsible for glutamate synthesis (Pinteaux et al., 1996); as well as the excitatory amino acid transporters that together with glutamine synthase are involved in the regulation of extraneuronal levels of glutamate (Volterra et al., 1994). Last, but not least, the NMDA receptor itself is highly sensitive to redox modulation through a redox-sensitive site (Herin and Aizenman, 2004), and decreases in the main antioxidant in brain, GSH, or reduced activity of GSH-peroxidase lead to oxidized hypofunctional NMDA-Rs (Jiang et al., 2000; Steullet et al., 2006). In particular, receptors composed of NR1:NR2A subunits were shown to have a highly reversible and rapid current potentiation by sulfhydryl redox agents, including GSH, acting on a specific redox site in NR2A (Kohr et al., 1994). The oxidation status of this redox site can affect the regulation of these receptors by spermine and protons, as well as the inhibition mediated by the high-affinity  $Zn^{2+}$  site (Lipton et al., 2002). On the other hand, oxidation of receptors containing NR2C is slowly reversible (Kohr et al., 1994). Given this heightened response to oxidative conditions, NMDA-R function in PV-interneurons may remain blocked when the IL-6/Nox2 pathway is activated, and this could lead to the enduring changes we observed in the phenotype of PV-interneurons in adult animals (Behrens et al., 2007, 2008). The less oxidizing conditions produced by inactivation of the IL-6/Nox2 pathway upon drug washout would then slowly reverse these effects.

Under normal physiological conditions the brain maintains a physiological range of superoxide production that is required for normal nervous system function. Perturbations of superoxide levels in either direction are associated with impaired LTP, altered glutamatergic neurotransmission, and poorer cognitive performance (Kishida and Klann, 2007). Thus, it could be assumed that the effects of transient activation of the IL-6/Nox2 pathway may be a consequence of normal regulatory mechanisms in brain, where brief activation of the pathway does not lead to enduring effects on inhibitory circuits. Indeed, this is what is observed 24 h after one injection of NMDA-R antagonists (Fig. 1a and Zuo et al., 2007; Behrens et al., 2008). Repetitive exposures in adulthood, however, through disinhibition-induced activation of the IL-6/Nox2 pathway induce an enduring, albeit reversible, dysfunction of the PV-interneuronal system (Fig. 1b).

Although the effects of activation of the IL-6/Nox2 pathway appear to be reversible in the adult brain, similar exposures during the second postnatal week produce an irreversible loss of PV-interneurons (Wang et al., 2008 and our unpublished results). GAD67 expression is required for the correct development of PV-synaptic contacts during postnatal development (Chattopadhyaya et al., 2007) and NMDA-R antagonists decrease the expression of this enzyme in PV-interneurons (Kinney et al., 2006; Behrens et al., 2007, 2008). Thus, it is possible that the decreased expression of GAD67 caused by NMDA-R antagonists during the second week of postnatal development could halt the maturation of PV-interneurons, profoundly affecting the development of this critical inhibitory system (Fig. 1c). This would impair the development of cortical networks involved in gamma-frequency generation and synchrony, and could eventually lead to the cognitive dysfunctions observed in schizophrenia. These effects should be more pronounced in at risk individuals that show diminished antioxidant defenses, as suggested for schizophrenia patients carrying specific single nucleotide polymorphisms in genes coding for the enzymes involved in GSH synthesis (reviewed in Do et al., 2009).



**Fig. 1.** Schematic representation of the effects of activation of the IL-6/Nox2 pathway on the PV-interneuronal system in the adult and perinatal brain. a) PV-interneuronal networks, through feedforward inhibition, control cortical output and generate oscillatory synchrony in the adult brain. b) In the adult brain, the initial disinhibition (increased glutamate) caused by repetitive exposures to sub-anesthetic concentrations of NMDA-R antagonists (i.e. ketamine) leads to the sustained activation of the IL-6/Nox2 pathway. The superoxide thus produced tips the redox balance in brain, leading to the reversible loss of GABAergic phenotype of PV-interneurons. c) If the activation of the IL-6/Nox2 pathway is triggered by repetitive exposure to NMDA-R antagonists during the maturation of PV-interneuronal networks, it produces irreversible loss of PV-interneurons and permanent dysfunction of the PV-interneuronal system in adulthood.

## 8. Summary

The evidence reviewed here points toward a precipitating oxidative period early in the development of PV-interneurons that, after a cascade of compensatory changes to other neurons, may leave the cortex in a highly vulnerable state. This may account for the long delay before the symptoms of schizophrenia appear in early adulthood, when synaptic reorganization, hormonal changes and environmental stresses could tip the balance toward the dysregulation of cortical circuits (Fig. 1). Only a few of the molecular mechanisms that might be responsible for this long chain of events have been uncovered so far. In particular the activation of Nox2 by IL-6 may trigger the oxidation of the NMDA-R on PV-interneurons during the critical period of their maturation, leading to the permanent

downregulation of GABAergic transmission by these interneurons. Critical experiments to test this hypothesis *in vivo* are underway.

## References

- Alexander, G.E., Goldman, P.S., 1978. Functional development of the dorsolateral prefrontal cortex: an analysis utilizing reversible cryogenic depression. *Brain Res.* 143, 233–249.
- Aoyama, K., Suh, S.W., Hamby, A.M., Liu, J., Chan, W.Y., Chen, Y., Swanson, R.A., 2006. Neuronal glutathione deficiency and age-dependent neurodegeneration in the EAAC1 deficient mouse. *Nat. Neurosci.* 9, 119–126.
- Ba, A., Seri, B.V., 1995. Psychomotor functions in developing rats: ontogenetic approach to structure-function relationships. *Neurosci. Biobehav. Rev.* 19, 413–425.
- Bachevalier, J., Beaugregard, M., 1993. Maturation of medial temporal lobe memory functions in rodents, monkeys, and humans. *Hippocampus* 3, 191–201.
- Bartos, M., Vida, I., Jonas, P., 2007. Synaptic mechanisms of synchronized gamma oscillations in inhibitory interneuron networks. *Nat. Rev. Neurosci.* 8, 45–56.
- Beasley, C.L., Reynolds, G.P., 1997. Parvalbumin-immunoreactive neurons are reduced in the prefrontal cortex of schizophrenics. *Schizophr. Res.* 24, 349–355.
- Behrens, M.M., Ali, S.S., Dao, D.N., Lucero, J., Shekhtman, G., Quick, K.L., Dugan, L.L., 2007. Ketamine-induced loss of phenotype of fast-spiking interneurons is mediated by NADPH-oxidase. *Science* 318, 1645–1647.
- Behrens, M.M., Ali, S.S., Dugan, L.L., 2008. Interleukin-6 mediates the increase in NADPH-oxidase in the ketamine model of schizophrenia. *J. Neurosci.* 28, 13957–13966.
- Belforte, J.E., Zsiros, V., Sklar, E.R., Yu, G., Li, Y., Quinlan, E.M., Nakazawa, K., 2008. Postnatal ablation of NMDA receptors in corticolimbic interneurons leads to schizophrenia-like behavior. *Soc. Neurosci. Abs.*
- Benes, F.M., Berretta, S., 2001. GABAergic interneurons: implications for understanding schizophrenia and bipolar disorder. *Neuropsychopharmacology* 25, 1–27.
- Berk, M., Copolov, D., Dean, O., Lu, K., Jeavons, S., Schapkaite, I., Anderson-Hunt, M., Judd, F., Katz, F., Katz, P., Ording-Jespersen, S., Little, J., Conus, P., Cuenod, M., Do, K.Q., Bush, A.I., 2008. *N*-Acetyl cysteine as a glutathione precursor for schizophrenia – a double-blind, randomized, placebo-controlled trial. *Biol. Psychiatry* 64, 361–368.
- Brown, A.S., 2006. Prenatal infection as a risk factor for schizophrenia. *Schizophr. Bull.* 32, 200–202.
- Butterfield, D.A., 2006. Oxidative stress in neurodegenerative disorders. *Antioxid. Redox Signal.* 8, 1971–1973.
- Cabungcal, J.H., Preissmann, D., Delseth, C., Cuenod, M., Do, K.Q., Schenk, F., 2007. Transitory glutathione deficit during brain development induces cognitive impairment in juvenile and adult rats: relevance to schizophrenia. *Neurobiol. Dis.* 26, 634–645.
- Cardin, J.A., Carlén, M., Meletis, K., Knoblich, U., Zhang, F., Deisseroth, K., Tsai, L.H., Moore, C.I., 2009. Driving fast-spiking cells induces gamma rhythm and controls sensory responses. *Nature* 459, 663–667.
- Cauli, B., Audinat, E., Lambolez, B., Angulo, M.C., Ropert, N., Tsuzuki, K., Hestrin, S., Rossier, J., 1997. Molecular and physiological diversity of cortical nonpyramidal cells. *J. Neurosci.* 17, 3894–3906.
- Chattopadhyaya, B., Di Cristo, G., Wu, C.Z., Knott, G., Kuhlman, S., Fu, Y., Palmiter, R.D., Huang, Z.J., 2007. GAD67-mediated GABA synthesis and signaling regulate inhibitory synaptic innervation in the visual cortex. *Neuron* 54 (6), 889–903.
- Cochran, S.M., Fujimura, M., Morris, B.J., Pratt, J.A., 2002. Acute and delayed effects of phencyclidine upon mRNA levels of markers of glutamatergic and GABAergic neurotransmitter function in the rat brain. *Synapse* 46, 206–214.
- Cochran, S.M., Kennedy, M., McKerchar, C.E., Steward, L.J., Pratt, J.A., Morris, B.J., 2003. Induction of metabolic hypofunction and neurochemical deficits after chronic intermittent exposure to phencyclidine: differential modulation by antipsychotic drugs. *Neuropsychopharmacology* 28, 265–275.
- Cruz, D.A., Eggan, S.M., Lewis, D.A., 2003. Postnatal development of pre- and postsynaptic GABA markers at chandelier cell connections with pyramidal neurons in monkey prefrontal cortex. *J. Comp. Neurol.* 465, 385–400.
- Dean, O., Bush, A.I., Berk, M., Copolov, D.L., van den Buuse, M., 2009. Glutathione depletion in the brain disrupts short-term spatial memory in the Y-maze in rats and mice. *Behav. Brain Res.* 198, 258–262.
- Di Cristo, G., Wu, C., Chattopadhyaya, B., Ango, F., Knott, G., Welker, E., Svoboda, K., Huang, Z.J., 2004. Subcellular domain-restricted GABAergic innervation in primary visual cortex in the absence of sensory and thalamic inputs. *Nat. Neurosci.* 7, 1184–1186.
- Do, K.Q., Cabungcal, J.H., Frank, A., Steullet, P., Cuenod, M., 2009. Redox dysregulation, neurodevelopment and schizophrenia. *Curr. Opin. Neurobiol.*, [Epub ahead of print].
- Do, K.Q., Trabesinger, A.H., Kirsten-Kruger, M., Lauer, C.J., Dydak, U., Hell, D., Holsboer, F., Boesiger, P., Cuenod, M., 2000. Schizophrenia: glutathione deficit in cerebrospinal fluid and prefrontal cortex *in vivo*. *Eur. J. Neurosci.* 12, 3721–3728.
- Doischer, D., Hosp, J.A., Yanagawa, Y., Obata, K., Jonas, P., Vida, I., Bartos, M., 2008. Postnatal differentiation of basket cells from slow to fast signaling devices. *J. Neurosci.* 28, 12956–12968.
- Feinberg, I., 1982. Schizophrenia: caused by a fault in programmed synaptic elimination during adolescence? *J. Psychiatr. Res.* 17 (4), 319–334.
- Fejgin, K., Palsson, E., Wass, C., Svensson, L., Klamer, D., 2008. Nitric oxide signaling in the medial prefrontal cortex is involved in the biochemical and behavioral effects of phencyclidine. *Neuropsychopharmacology* 33, 1874–1883.
- Gainetdinov, R.R., Mohn, A.R., Caron, M.G., 2001. Genetic animal models: focus on schizophrenia. *Trends Neurosci.* 24, 527–533.
- Ganguli, R., Yang, Z., Shurin, G., Chengappa, K.N., Brar, J.S., Gubbi, A.V., Rabin, B.S., 1994. Serum interleukin-6 concentration in schizophrenia: elevation associated with duration of illness. *Psychiatry Res.* 51, 1–10.
- Garrity, A.G., Pearlson, G.D., McKiernan, K., Lloyd, D., Kiehl, K.A., Calhoun, V.D., 2007. Aberrant “default mode” functional connectivity in schizophrenia. *Am. J. Psychiatry* 164, 450–457.
- Gilmore, J.H., Jarskog, L.F., 1997. Exposure to infection and brain development: cytokines in the pathogenesis of schizophrenia. *Schizophr. Res.* 24, 365–367.
- Goldberg, J.H., Yuste, R., Tamas, G., 2003.  $Ca^{2+}$  imaging of mouse neocortical interneurone dendrites: contribution of  $Ca^{2+}$ -permeable AMPA and NMDA receptors to subthreshold  $Ca^{2+}$  dynamics. *J. Physiol.* 551, 67–78.
- Goldman-Rakic, P.S., 1999. The physiological approach: functional architecture of working memory and disordered cognition in schizophrenia. *Biol. Psychiatry* 46, 650–661.
- Gonzalez-Burgos, G., Lewis, D.A., 2008. GABA neurons and the mechanisms of network oscillations: implications for understanding cortical dysfunction in schizophrenia. *Schizophr. Bull.* 34, 944–961.
- Grunze, H.C., Rainnie, D.G., Hasselmo, M.E., Barkai, E., Hearn, E.F., McCarley, R.W., Greene, R.W., 1996. NMDA-dependent modulation of CA1 local circuit inhibition. *J. Neurosci.* 16, 2034–2043.
- Gysin, R., Kraftsik, R., Sandell, J., Bovet, P., Chappuis, C., Conus, P., Deppen, P., Preisig, M., Ruiz, V., Steullet, P., Tosic, M., Werge, T., Cuenod, M., Do, K.Q., 2007. Impaired glutathione synthesis in schizophrenia: convergent genetic and functional evidence. *Proc. Natl. Acad. Sci. U.S.A.* 104, 16621–16626.
- Hashimoto, T., Volk, D.W., Eggan, S.M., Mirmics, K., Pierri, J.N., Sun, Z., Sampson, A.R., Lewis, D.A., 2003. Gene expression deficits in a subclass of GABA neurons in the prefrontal cortex of subjects with schizophrenia. *J. Neurosci.* 23, 6315–6326.
- Herin, G.A., Aizenman, E., 2004. Amino terminal domain regulation of NMDA receptor function. *Eur. J. Pharmacol.* 500, 101–111.
- Hikida, T., Jaaro-Peled, H., Seshadri, S., Oishi, K., Hookway, C., Kong, S., Wu, D., Xue, R., Andrade, M., Tankou, S., Mori, S., Gallagher, M., Ishizuka, K., Pletnikov, M., Kida, S., Sawada, A., 2007. Dominant-negative DISC1 transgenic mice display schizophrenia-associated phenotypes detected by measures translatable to humans. *Proc. Natl. Acad. Sci. U.S.A.* 104, 1451–14506.
- Holcomb, H.H., Lahti, A.C., Medoff, D.R., Cullen, T., Tamminga, C.A., 2005. Effects of noncompetitive NMDA receptor blockade on anterior cingulate cerebral blood flow in volunteers with schizophrenia. *Neuropsychopharmacology* 30, 2275–2282.
- Homayoun, H., Moghaddam, B., 2007. NMDA receptor hypofunction produces opposite effects on prefrontal cortex interneurons and pyramidal neurons. *J. Neurosci.* 27, 11496–11500.
- van Hoof, J.A., Giuffrida, R., Blatow, M., Monyer, H., 2000. Differential expression of group I metabotropic glutamate receptors in functionally distinct hippocampal interneurons. *J. Neurosci.* 20, 3544–3551.
- Huang, Z.J., 2009. Activity-dependent development of inhibitory synapses and innervation pattern: role of GABA signalling and beyond. *J. Physiol.* 587, 1881–1888.
- Jacobsen, J.P., Rodriguiz, R.M., Mork, A., Wetsel, W.C., 2005. Monoaminergic dysregulation in glutathione-deficient mice: possible relevance to schizophrenia? *Neuroscience* 132, 1055–1072.
- Jentsch, J.D., Roth, R.H., 1999. The neuropsychopharmacology of phencyclidine: from NMDA receptor hypofunction to the dopamine hypothesis of schizophrenia. *Neuropsychopharmacology* 20, 201–225.
- Jiang, D., Akopian, G., Ho, Y.S., Walsh, J.P., Andersen, J.K., 2000. Chronic brain oxidation in a glutathione peroxidase knockout mouse model results in increased resistance to induced epileptic seizures. *Exp. Neurol.* 164, 257–268.
- Kawaguchi, Y., Kubota, Y., 1993. Correlation of physiological subgroupings of non-pyramidal cells with parvalbumin- and calbindinD28k-immunoreactive neurons in layer V of rat frontal cortex. *J. Neurophysiol.* 70, 387–396.
- Keilhoff, G., Becker, A., Grecksch, G., Wolf, G., Bernstein, H.G., 2004. Repeated application of ketamine to rats induces changes in the hippocampal expression of parvalbumin, neuronal nitric oxide synthase and cFOS similar to those found in human schizophrenia. *Neuroscience* 126, 591–598.
- Kerns, J.G., Nuechterlein, K.H., Braver, T.S., Barch, D.M., 2008. Executive functioning component mechanisms and schizophrenia. *Biol. Psychiatry* 64, 26–33.
- Kinney, J.W., Davis, C.N., Tabarean, I., Conti, B., Bartfai, T., Behrens, M.M., 2006. A specific role for NR2A-containing NMDA receptors in the maintenance of parvalbumin and GAD67 immunoreactivity in cultured interneurons. *J. Neurosci.* 26, 1604–1615.
- Kishida, K.T., Klann, E., 2007. Sources and targets of reactive oxygen species in synaptic plasticity and memory. *Antioxid. Redox Signal.* 9, 233–244.
- Kohr, G., Eckardt, S., Luddens, H., Monyer, H., Seeburg, P.H., 1994. NMDA receptor channels: subunit-specific potentiation by reducing agents. *Neuron* 12, 1031–1040.
- Kotermanski, S.E., Johnson, J.W., 2009.  $Mg^{2+}$  imparts NMDA receptor subtype selectivity to the Alzheimer’s drug memantine. *J. Neurosci.* 29, 2774–2779.
- Krystal, J.H., Perry Jr., E.B., Gueorguieva, R., Belger, A., Madonick, S.H., Abi-Dargham, A., Cooper, T.B., MacDougall, L., Abi-Saab, W., D’Souza, D.C., 2005. Comparative and interactive human psychopharmacologic effects of ketamine and amphetamine: implications for glutamatergic and dopaminergic model psychoses and cognitive function. *Arch. Gen. Psychiatry* 62, 985–994.

- Kudoh, A., Takase, H., Takahira, Y., Katagai, H., Takazawa, T., 2003. Postoperative confusion in schizophrenic patients is affected by interleukin-6. *J. Clin. Anesth.* 15, 455–462.
- Laruelle, M., Abi-Dargham, A., Casanova, M.F., Toti, R., Weinberger, D.R., Kleinman, J.E., 1993. Selective abnormalities of prefrontal serotonergic receptors in schizophrenia. A postmortem study. *Arch. Gen. Psychiatry* 50, 810–818.
- Lavoie, S., Murray, M.M., Deppen, P., Knyazeva, M.G., Berk, M., Boulat, O., Bovet, P., Bush, A.L., Conus, P., Copolov, D., Fornari, E., Meuli, R., Solida, A., Vianin, P., Cuenod, M., Buclin, T., Do, K.Q., 2008. Glutathione precursor, N-acetyl-cysteine, improves mismatch negativity in schizophrenia patients. *Neuropsychopharmacology* 33, 2187–2199.
- Lewis, D.A., Cruz, D., Eggen, S., Erickson, S., 2004. Postnatal development of prefrontal inhibitory circuits and the pathophysiology of cognitive dysfunction in schizophrenia. *Ann. N. Y. Acad. Sci.* 1021, 64–76.
- Lewis, D.A., Hashimoto, T., Volk, D.W., 2005. Cortical inhibitory neurons and schizophrenia. *Nat. Rev. Neurosci.* 6, 312–324.
- Li, Q., Clark, S., Lewis, D.V., Wilson, W.A., 2002. NMDA receptor antagonists disinhibit rat posterior cingulate and retrosplenial cortices: a potential mechanism of neurotoxicity. *J. Neurosci.* 22, 3070–3080.
- Lin, A., Kenis, G., Bignotti, S., Tura, G.J., De Jong, R., Bosmans, E., Pioli, R., Altamura, C., Scharpe, S., Maes, M., 1998. The inflammatory response system in treatment-resistant schizophrenia: increased serum interleukin-6. *Schizophr. Res.* 32, 9–15.
- Lipton, S.A., Choi, Y.B., Takahashi, H., Zhang, D., Li, W., Godzik, A., Bankston, L.A., 2002. Cysteine regulation of protein function – as exemplified by NMDA-receptor modulation. *Trends Neurosci.* 25, 474–480.
- Lisman, J.E., Coyle, J.T., Green, R.W., Javitt, D.C., Benes, F.M., Heckers, S., Grace, A.A., 2008. Circuit-based framework for understanding neurotransmitter and risk gene interactions in schizophrenia. *Trends Neurosci.* 31, 234–242.
- Lodge, D.J., Behrens, M.M., Grace, A.A., 2009. A loss of parvalbumin-containing interneurons is associated with diminished oscillatory activity in an animal model of schizophrenia. *J. Neurosci.* 29, 2344–2354.
- Lodge, D.J., Grace, A.A., 2007. Aberrant hippocampal activity underlies the dopamine dysregulation in an animal model of schizophrenia. *J. Neurosci.* 27, 11424–11430.
- Markram, H., Toledo-Rodriguez, M., Wang, Y., Gupta, A., Silberberg, G., Wu, C., 2004. Interneurons of the neocortical inhibitory system. *Nat. Rev. Neurosci.* 5, 793–807.
- Mellon, R.D., Simone, A.F., Rappaport, B.A., 2007. Use of anesthetic agents in neonates and young children. *Anesth. Analg.* 104, 509–520.
- Meyer, U., Nyffeler, M., Yee, B.K., Knuesel, I., Feldon, J., 2007. Adult brain and behavioral pathological markers of prenatal immune challenge during early/middle and late fetal development in mice. *Brain Behav. Immun.* 22, 469–486.
- Middleton, S., Jalics, J., Kispersky, T., Lebeau, F.E., Roopun, A.K., Kopell, N.J., Whittington, M.A., Cunningham, M.O., 2008. NMDA receptor-dependent switching between different gamma rhythm-generating microcircuits in entorhinal cortex. *Proc. Natl. Acad. Sci. U.S.A.* 105, 18572–18577.
- Morris, B.J., Cochran, S.M., Pratt, J.A., 2005. PCP: from pharmacology to modelling schizophrenia. *Curr. Opin. Pharmacol.* 5, 101–106.
- Morrow, B.A., Elsworth, J.D., Roth, R.H., 2007. Repeated phencyclidine in monkeys results in loss of parvalbumin-containing axo-axonic projections in the prefrontal cortex. *Psychopharmacology (Berl.)* 192, 283–290.
- Mouri, A., Noda, Y., Enomoto, T., Nabeshima, T., 2007. Phencyclidine animal models of schizophrenia: approaches from abnormality of glutamatergic neurotransmission and neurodevelopment. *Neurochem. Int.* 51, 173–184.
- Muller, N., Riedel, M., Gruber, R., Ackenheil, M., Schwarz, M.J., 2000. The immune system and schizophrenia. An integrative view. *Ann. N. Y. Acad. Sci.* 917, 456–467.
- Mustafa, A.K., Kumar, M., Selvakumar, B., Ho, G.P., Ehmsen, J.T., Barrow, R.K., Amzel, L.M., Snyder, S.H., 2007. Nitric oxide S-nitrosylates serine racemase, mediating feedback inhibition of D-serine formation. *Proc. Natl. Acad. Sci. U.S.A.* 104, 2950–2955.
- Naudin, J., Mege, J.L., Azorin, J.M., Dassa, D., 1996. Elevated circulating levels of IL-6 in schizophrenia. *Schizophr. Res.* 20, 269–273.
- Nawa, H., Takahashi, M., Patterson, P.H., 2000. Cytokine and growth factor involvement in schizophrenia – support for the developmental model. *Mol. Psychiatry* 5, 594–603.
- Nudmamud, S., Reynolds, L.M., Reynolds, G.P., 2003. N-acetylaspartate and N-acetylaspartylglutamate deficits in superior temporal cortex in schizophrenia and bipolar disorder: a postmortem study. *Biol. Psychiatry* 53, 1138–1141.
- Nunes, S.O., Borelli, S.D., Matsuo, T., Watanabe, M.A., Itano, E.N., 2005. The association of the HLA in patients with schizophrenia, schizoaffective disorder, and in their biological relatives. *Schizophr. Res.* 76, 195–198.
- Okaty, B.W., Miller, M.N., Sugino, K., Hempel, C.M., Nelson, S.B., 2009. Transcriptional and electrophysiological maturation of neocortical fast-spiking GABAergic interneurons. *J. Neurosci.* 29, 7040–7052.
- Olney, J.W., 2002. New insights and new issues in developmental neurotoxicology. *Neurotoxicology* 23, 659–668.
- Olney, J.W., Newcomer, J.W., Farber, N.B., 1999. NMDA receptor hypofunction model of schizophrenia. *J. Psychiatr. Res.* 33, 523–533.
- Papadia, S., Soriano, F.X., Leveille, F., Martel, M.A., Dakin, K.A., Hansen, H.H., Kaindl, A., Siffringer, M., Fowler, J., Stefovská, V., McKenzie, G., Craigm, M., Corriveau, R., Ghazal, P., Horsburgh, K., Yankner, B.A., Wyllie, D.J., Ikonomidou, C., Hardingham, G.E., 2008. Synaptic NMDA receptor activity boosts intrinsic antioxidant defenses. *Nat. Neurosci.* 11, 476–487.
- Patterson, P.H., 2008. Immune involvement in schizophrenia and autism: etiology, pathology and animal models. *Behav. Brain Res.* [Epub ahead of print].
- Pileblad, E., Magnusson, T., Fornstedt, B., 1989. Reduction of brain glutathione by L-buthionine sulfoximine potentiates the dopamine-depleting action of 6-hydroxydopamine in rat striatum. *J. Neurochem.* 52, 978–980.
- Pinteaux, E., Copin, J.C., Ledig, M., Tholey, G., 1996. Modulation of oxygen-radical-scavenging enzymes by oxidative stress in primary cultures of rat astroglial cells. *Dev. Neurosci.* 18, 397–404.
- Pollmacher, T., Haack, M., Schuld, A., Kraus, T., Hinze-Selch, D., 2000. Effects of antipsychotic drugs on cytokine networks. *J. Psychiatr. Res.* 34, 369–382.
- Rao, S.G., Williams, G.V., Goldman-Rakic, P.S., 2000. Destruction and creation of spatial tuning by disinhibition: GABA(A) blockade of prefrontal cortical neurons engaged by working memory. *J. Neurosci.* 20, 485–494.
- Rapoport, J.L., Addington, A.M., Frangou, S., Psych, M.R., 2005. The neurodevelopmental model of schizophrenia: update 2005. *Mol. Psychiatry* 10, 434–449.
- Reynolds, G.P., Beasley, C.L., 2001. GABAergic neuronal subtypes in the human frontal cortex – development and deficits in schizophrenia. *J. Chem. Neuroanat.* 22, 95–100.
- Reynolds, L.M., Cochran, S.M., Morris, B.J., Pratt, J.A., Reynolds, G.P., 2005. Chronic phencyclidine administration induces schizophrenia-like changes in N-acetylaspartate and N-acetylaspartylglutamate in rat brain. *Schizophr. Res.* 73, 147–152.
- Roopun, A.K., Cunningham, M.O., Racca, C., Alter, K., Traub, R.D., Whittington, M.A., 2008. Region-specific changes in gamma and beta2 rhythms in NMDA receptor dysfunction models of schizophrenia. *Schizophr. Bull.* 34, 962–973.
- Rowan, M.J., Klyubin, I., Wang, Q., Anwyl, R., 2005. Synaptic plasticity disruption by amyloid beta protein: modulation by potential Alzheimer's disease modifying therapies. *Biochem. Soc. Trans.* 33, 563–567.
- Rujescu, D., Bender, A., Keck, M., Hartmann, A.M., Ohl, F., Raeder, H., Giegling, I., Genius, J., McCarley, R.V., Moller, H.J., Grunze, H., 2006. A pharmacological model for psychosis based on N-methyl-D-aspartate receptor hypofunction: molecular, cellular, functional and behavioral abnormalities. *Biol. Psychiatry* 59, 721–729.
- Salinas, E., Sejnowski, T.J., 2001. Correlated neuronal activity and the flow of neural information. *Nat. Rev. Neurosci.* 2, 539–550.
- Satoh, T., Lipton, S.A., 2007. Redox regulation of neuronal survival mediated by electrophilic compounds. *Trends Neurosci.* 30, 37–45.
- Sirota, P., Gavrieli, R., Wolach, B., 2003. Overproduction of neutrophil radical oxygen species correlates with negative symptoms in schizophrenic patients: parallel studies on neutrophil chemotaxis, superoxide production and bactericidal activity. *Psychiatry Res.* 121, 123–132.
- Smith, S.E., Li, J., Garbett, K., Mirmics, K., Patterson, P.H., 2007. Maternal immune activation alters fetal brain development through interleukin-6. *J. Neurosci.* 27, 10695–10702.
- Sohal, V.S., Zhang, F., Yizhar, O., Deisseroth, K., 2009. Parvalbumin neurons and gamma rhythms enhance cortical circuit performance. *Nature* 459, 698–702.
- Stefani, M.R., Moghaddam, B., 2005a. Transient N-methyl-D-aspartate receptor blockade in early development causes lasting cognitive deficits relevant to schizophrenia. *Biol. Psychiatry* 57, 433–436.
- Stefani, M.R., Moghaddam, B., 2005b. Systemic and prefrontal cortical NMDA receptor blockade differentially affect discrimination learning and set-shift ability in rats. *Behav. Neurosci.* 119, 420–428.
- Steulet, P., Neijt, H.C., Cuenod, M., Do, K.Q., 2006. Synaptic plasticity impairment and hypofunction of NMDA receptors induced by glutathione deficit: relevance to schizophrenia. *Neuroscience* 137, 807–819.
- Steward, L.J., Kennedy, M.D., Morris, B.J., Pratt, J.A., 2004. The atypical antipsychotic drug clozapine enhances chronic PCP-induced regulation of prefrontal cortex 5-HT2A receptors. *Neuropharmacology* 47, 527–537.
- Stoet, G., Snyder, L.H., 2006. Effects of the NMDA antagonist ketamine on task-switching performance: evidence for specific impairments of executive control. *Neuropsychopharmacology* 31, 1675–1681.
- Takahata, R., Moghaddam, B., 2003. Activation of glutamate neurotransmission in the prefrontal cortex sustains the motoric and dopaminergic effects of phencyclidine. *Neuropsychopharmacology* 28, 1117–1124.
- Tomitaka, S., Tomitaka, M., Tolliver, B.K., Sharp, F.R., 2000. Bilateral blockade of NMDA receptors in anterior thalamus by dizocilpine (MK-801) injures pyramidal neurons in rat retrosplenial cortex. *Eur. J. Neurosci.* 12, 1420–1430.
- Tosic, M., Ott, J., Barral, S., Bovet, P., Deppen, P., Gheorghita, F., Matthey, M.L., Parnas, J., Preisig, M., Saraga, M., Solida, A., Timm, S., Wang, A.G., Werge, T., Cuenod, M., Do, K.Q., 2006. Schizophrenia and oxidative stress: glutamate cysteine ligase modifier as a susceptibility gene. *Am. J. Hum. Genet.* 79, 586–592.
- Trotti, D., Rizzini, B.L., Rossi, D., Haugeto, O., Racagni, G., Danbolt, N.C., Volterra, A., 1997. Neuronal and glial glutamate transporters possess an SH-based redox regulatory mechanism. *Eur. J. Neurosci.* 9, 1236–1243.
- Tseng, K.Y., O'Donnell, P., 2007. Dopamine modulation of prefrontal cortical interneurons changes during adolescence. *Cereb. Cortex* 17, 1235–1240.
- Uhlhaas, P.J., Roux, F., Singer, W., Haenschel, C., Sireteanu, R., Rodriguez, E., 2009. The development of neural synchrony reflects late maturation and restructuring of functional networks in humans. *Natl. Acad. Sci. U.S.A.* 106, 9866–9871.
- Uhlhaas, P.J., Haenschel, C., Nikolic, D., Singer, W., 2008. The role of oscillations and synchrony in cortical networks and their putative relevance for the pathophysiology of schizophrenia. *Schizophr. Bull.* 34, 927–943.
- Volterra, A., Trotti, D., Floridi, S., Racagni, G., 1994. Reactive oxygen species inhibit high-affinity glutamate uptake: a molecular mechanism and neuropathological implications. *Ann. N. Y. Acad. Sci.* 738, 153–162.
- Wang, C., McInnis, J., West, J.B., Bao, J., Anastasio, N., Guidry, J.A., Ye, Y., Salvemini, D., Johnson, K.M., 2003. Blockade of phencyclidine-induced cortical apoptosis and

- deficits in prepulse inhibition by M40403, a superoxide dismutase mimetic. *J. Pharmacol. Exp. Ther.* 304, 266–271.
- Wang, C.Z., Yang, S.F., Xia, Y., Johnson, K.M., 2008. Postnatal phencyclidine administration selectively reduces adult cortical parvalbumin-containing interneurons. *Neuropsychopharmacology* 33, 2442–2455.
- Wang, C., Slikker Jr., W., 2008. Strategies and experimental models for evaluating anesthetics: effects on the developing nervous system. *Anesth. Analg.* 106, 1643–1658.
- Wilson, F.A., O'Scalaidhe, S.P., Goldman-Rakic, P.S., 1994. Functional synergism between putative gamma-aminobutyrate-containing neurons and pyramidal neurons in prefrontal cortex. *Proc. Natl. Acad. Sci. U.S.A.* 91, 4009–4013.
- Xi, D., Keeler, B., Zhang, W., Houle, J.D., Gao, W.J., 2009. NMDA receptor subunit expression in GABAergic interneurons in the prefrontal cortex: application of laser microdissection technique. *J. Neurosci. Methods* 176, 172–181.
- Xia, S., Cai, Z.Y., Thi, L.L., Kim-Han, J.S., Dugan, L.L., Covey, D.F., Rothman, S.M., 2002. The estrogen receptor is not essential for all estrogen neuroprotection: new evidence from a new analog. *Neurobiol. Dis.* 9, 282–293.
- Yao, J.K., Leonard, S., Reddy, R., 2006. Altered glutathione redox state in schizophrenia. *Dis. Markers* 22, 83–93.
- Zhang, X.Y., Zhou, D.F., Zhang, P.Y., Wu, G.Y., Cao, L.Y., Shen, Y.C., 2002. Elevated interleukin-2, interleukin-6 and interleukin-8 serum levels in neuroleptic-free schizophrenia: association with psychopathology. *Schizophr. Res.* 57, 247–258.
- Zhang, Y., Behrens, M.M., Lisman, J.E., 2008. Prolonged exposure to NMDAR antagonist suppresses inhibitory synaptic transmission in prefrontal cortex. *J. Neurophysiol.* 100, 959–965.
- Zuo, D.Y., Wu, Y.L., Yao, W.X., Cao, Y., Wu, C.F., Tanaka, M., 2007. Effect of MK-801 and ketamine on hydroxyl radical generation in the posterior cingulate and retrosplenial cortex of free-moving mice, as determined by in vivo microdialysis. *Pharmacol. Biochem. Behav.* 86, 1–7.